

An estimate for the volume entropy of nonpositively curved graph manifolds

S. Buyalo *

Abstract

Let M be a closed 3-dimensional graph manifold. We prove that $h(g) > 1$ for each geometrization g of M , where $h(g)$ is the topological entropy of geodesic flow of g .

1 Introduction

Asymptotic geometry of a nonpositively curved (NPC for brevity) graph manifold is a complicated mixture of flat and hyperbolic parts which both nontrivially contribute in general picture. Recall that a NPC metric on a closed 3-dimensional graph manifold M recovers the JSJ-decomposition of M in the following sense. There is the unique (up to isotopy) minimal finite collection E of flat geodesically embedded tori and Klein bottles which are pairwise disjoint, and the metric completion of each connected component of the complement to E is a Seifert space called a block of M . Each block M_v is fibered over a 2-orbifold S_v with negative Euler characteristic, $\chi(S_v) < 0$. Furthermore, the metric locally splits as $U \times (-\varepsilon, \varepsilon)$ along the interior of each block, where U is a NPC surface, the splitting is compatible with the fibration, the fibers are closed geodesics, and the regular fibers have one and the same length $l_v > 0$ depending only on the block.

Since we are interested in asymptotic properties, which are certainly the same for any finite covering of M , we may assume for simplicity that M is orientable, the collection E consists of tori, and each block is a trivial S^1 -bundle over a compact surface S_v with boundary, $M_v = S_v \times S^1$. We also assume that the graph manifold structure of M is nontrivial, i.e. M itself is not a Seifert fibered space (though it may consist of one block).

The flat part of the asymptotic geometry of M studied in [BS], [CK], see also [HS]. Roughly speaking, it can be described by very special geodesic rays $[0, \infty) \rightarrow M$, which terminate in no block, skip through separating tori $e \in E$ almost tangentially spending most of the time near tori and moreover this time rapidly increases with each step. Though the set of such rays is

*Partially supported by RFFI Grants 99-01-00104 and 00-15-96024.

a negligible part all of the rays, it contains an important information about geometry of M : it was shown in [CK] how this information allows to recover (up to scaling) the marked length spectrum of closed geodesics on S_v and the fiber length for each block M_v .

Here we study the hyperbolic part of the asymptotic geometry of M , assuming that the surface U from the local splitting $U \times (-\varepsilon, \varepsilon)$ above has the constant curvature $K = -1$. In other words, each block fibers over a *hyperbolic* orbifold (surface) S_v . A NPC metric on M satisfying this condition is called a *geometrization* of M (because the metric of each block is modelled on $H^2 \times \mathbb{R}$). Note that any geometrization of M is only $C^{1,1}$ -smooth being analytic along the interior of each block. It is known [L] that M admits a NPC metric if and only if it admits a geometrization. Necessary and sufficient topological conditions for M to carry a NPC metric were found in [BK].

The relevant metric invariant which measures hyperbolicity of a space is the volume entropy h . Let $\pi : X \rightarrow M$ be the universal covering, $x_0 \in X$. Recall that $h = h(X)$ is defined by

$$h = \lim_{R \rightarrow \infty} \frac{1}{R} \ln \text{vol } B_R(x_0),$$

where $B_R(x)$ is the ball in X of radius R centered at x . It is well known [M] that the limit exists, h is independent of the choice of x_0 and if M is NPC then h coincides with the topological entropy of geodesic flow of M . The volume entropy scales as l^{-1} , l is the length. Thus the choice of a geometrization g of M also serves as a normalization. The sectional curvatures of g satisfy $-1 \leq K \leq 0$. Hence $h(g) \leq 2$ by comparison with H^3 . Our main result is this.

Theorem 1.1. *For any geometrization g of a graph manifold M we have $h(g) > 1$.*

Remark 1.2. Though the universal covering X of M looks much more complicated than the model space $H^2 \times \mathbb{R}$, even the estimate $h(g) \geq 1 = h(H^2 \times \mathbb{R})$ is not obvious and nontrivial: X contains no isometrically and geodesically embedded H^2 , which would lead to $h(g) \geq 1$; on the other hand, the attempt to compare $X \rightarrow H^2 \times \mathbb{R}$ via exponential maps identifying some tangent spaces fails because the Jacobian of this map is > 1 at some points. Finally, the estimate from [BW] for the measure theoretic entropy of geodesic flow, which is always $\leq h(g)$, gives only $\pi/4 < 1$ as a lower bound for any geometrization g of M (if one ignores the fact that $C^{1,1}$ -smoothness of g is not sufficient to apply this estimate).

To prove Theorem 1.1 we use the well known fact that $h(g)$ coincides with the critical exponent of a Poincaré series

$$\mathcal{P}(t) = \sum_{\gamma \in \Gamma} e^{-t|x_0 - \gamma x_0|},$$

where the fundamental group $\Gamma = \pi_1(M)$ isometrically acts on X as the deck transformation group. Actually, instead of $\mathcal{P}$ we use a modified Poincaré series $\mathcal{P}_W$, where summation is taken over a set W of walls in X . Our proof involves three ingredients: (i) a local estimate, which is technical and used in (ii); this estimate is obtained in section 2. (ii) An accumulating procedure, which is an inductive construction of appropriately chosen broken geodesics in X between the base point x_0 and the walls from W ; the choice of these paths is the key point of the proof. The outcome of the accumulating procedure is a generating set for $\mathcal{P}_W$ used in (iii); the procedure is described in sect. 3. (iii) A self-similarity type argument. This part of the proof uses a standard idea from self-similarity theory to show that $\mathcal{P}_W(h)$ diverges for some $h > 1$ using the generating set obtained in (ii). This is done in sect. 4.

Acknowledgment. The author is grateful to W. Ballmann for useful discussions of the topic of this note.

2 Local estimate

Let F be the universal covering of a compact hyperbolic surface S with geodesic boundary. We identify F with a convex subset $F \subset \mathbb{H}^2$ bounded by countably many disjoint geodesic line and fix $o \in \mathbb{H}^2 \setminus F$. Let w_0 be the boundary line of F closest to o , $o_0 \in w_0$ be the point on w_0 closest to o , so that $|o - o_0| = \text{dist}(o, F) =: l > 0$.

Let A be the set of the boundary lines of F different from w_0 . For each $w \in A$ we denote by $o_w \in w$ the point closest to o . Then the geodesic segment oo_w intersects w_0 at some point t_w , and for $\tilde{l}_w = |o - o_w|$ we have

$$\tilde{l}_w = \tilde{l}'_w + l''_w,$$

where $\tilde{l}'_w = |o - t_w|$, $l''_w = |t_w - o_w|$ (all distances are taken in $\mathbb{H}^2$).

Next, we identify $\mathbb{H}^2$ with $\mathbb{H}^2 \times 0 \subset \mathbb{H}^2 \times \mathbb{R}$, so that F becomes a subset of $\mathbb{H}^2 \times \mathbb{R}$, and we use for it the notations introduced above. Note that the point o_0 is closest to o from the wall $w_0 \times \mathbb{R}$. We take a geodesic line $\sigma \subset w_0 \times \mathbb{R}$ through o_0 which is not horizontal, i.e., $\sigma \neq w_0 \times 0$, and take $s_w \in \sigma$ with $|s_w - o_0| = |t_w - o_0|$. Now, we put $l'_w := |o - s_w|$ (the distance is taken in $\mathbb{H}^2 \times \mathbb{R}$), $\Delta_w := \tilde{l}'_w - l'_w$.

In other words, we replace the distance $\tilde{l}'_w$ between o and t_w in the hyperbolic plane $\mathbb{H}^2$ by the distance l'_w in $\mathbb{H}^2 \times \mathbb{R}$, which is shorter by comparison: the triangles $oo_0t_w \subset \mathbb{H}^2 \times 0$, $oo_0s_w \subset \mathbb{H}^2 \times \mathbb{R}$ both have the right angles at o_0 , $\angle(oo_0t_w) = \frac{\pi}{2} = \angle(oo_0s_w)$, the common side oo_0 and equal sides $|o_0 - t_w| = |o_0 - s_w|$. Since oo_0t_w lies in the hyperbolic plane $\mathbb{H}^2 \times 0$ but oo_0s_w not, we have $\Delta_w > 0$ except the case $t_w = o_0 = s_w$. Now, we want to estimate accumulation of the differences Δ_w from below. The precise statement is this.

Lemma 2.1. *Given $l_0 > 0$, $\alpha_0 \in (0, \pi/2]$, there exists $\lambda_0 > 1$, which depends only on l_0 , α_0 and the compact surface S , so that*

$$\lambda(F, l, \alpha) := e^l \sum_{w \in A} e^{\Delta_w} e^{-\tilde{l}_w} \geq \lambda_0,$$

whenever $l = \text{dist}(o, w_0) \geq l_0$ and the angle α between the lines $w_0 \times 0$ and σ is at least α_0 , $\alpha_0 \leq \alpha \leq \pi/2$.

Proof. By well known formula of hyperbolic geometry we have $e^l = (\tan \frac{\psi}{4})^{-1}$, $e^{-\tilde{l}_w} = \tan \frac{\psi_w}{4}$, where ψ , ψ_w are the angles under which w_0 respectively $w \in A$ are observed in $\mathbb{H}^2$ from o . The boundary at infinity $\partial_\infty F \subset \partial_\infty \mathbb{H}^2 = S^1$ coincides with the limit set of $\pi_1(S)$ represented in $\text{Iso}(\mathbb{H}^2)$ as a Fuchsian group of second kind. It is well known that the Hausdorff dimension of $\partial_\infty F$ (with respect to the angle metric) is < 1 , in particular, the Lebesgue measure of $\partial_\infty F$ is zero. Thus $\psi = \sum_{w \in A} \psi_w$. Therefore, $\tan \frac{\psi}{4} \leq \sum_{w \in A} \tan \frac{\psi_w}{4}$ and

$$e^l \sum_{w \in A} e^{-\tilde{l}_w} = \sum_{w \in A} \tau_w \geq 1,$$

where $\tau_w = e^{l-\tilde{l}_w}$. However, this sum, $\sum_{w \in A} \tau_w$, can be close to 1 as much as we like (taking for instance $l \rightarrow \infty$). As we noticed above, $\Delta_w > 0$ unless $s_w = t_w$. So we always have $\lambda(F, l, \alpha) > 1$. The point is that $\lambda(F, l, \alpha)$ is separated from 1 uniformly over all $l \geq l_0$, $\alpha \geq \alpha_0$.

Consider $A_0 \subset A$ consisting of all $w \in A$ with $|t_w - o_0| \geq 1$. Then $\Delta_w \geq \delta_0 > 0$ for all $w \in A_0$, where δ_0 depends only on l_0 , α_0 . We claim that

$$\sum_{w \in A_0} \tau_w \geq m_0 > 0, \tag{1}$$

where $m_0 = m_0(\tilde{F}, l_0)$ is independent of l .

Assuming (1), we have

$$\begin{aligned} \lambda(F, l, \alpha) &= \sum_{w \in A} e^{\Delta_w} \tau_w \geq \sum_{w \in A_0} e^{\Delta_w} \tau_w + \sum_{w \in A \setminus A_0} \tau_w \\ &\geq e^{\delta_0} \sum_{w \in A_0} \tau_w + \sum_{w \in A \setminus A_0} \tau_w \\ &= (e^{\delta_0} - 1) \sum_{w \in A_0} \tau_w + \sum_{w \in A} \tau_w \geq (e^{\delta_0} - 1)m_0 + 1 =: \lambda_0 > 1. \end{aligned}$$

It remains to prove (1). Let $o'_w \in w$ be the point closest to $o_0 \in w_0$. Then $\tilde{l}_w \leq |o - o'_w| \leq l + |o_0 - o'_w|$, thus $\tilde{l}_w - l \leq \text{dist}(o_0, w)$. Hence $\tau_w \geq e^{-\text{dist}(o_0, w)} \geq \frac{\psi_w}{4}$, where w is observed from o_0 under the angle $\tilde{\psi}_w$.

The Lebesgue measure class on $\partial_\infty \mathbb{H}^2$ is independent of the choice of origin, thus $\sum_{w \in A} \tilde{\psi}_w = \pi$.

Since $l \geq l_0$, for a sufficiently small $m_0 = m_0(S, l_0) > 0$ the sectors $S^+(m_0)$, $S^-(m_0)$, defined below, intersect no $w \in A \setminus A_0$. Here is the definition of $S^\pm(m_0)$. The common vertex o_0 of $S^\pm(m_0)$ divides the line w_0 into two opposite rays $w_0^\pm$. The sectors $S^\pm(m_0) \subset \mathbb{H}^2$ are bounded by the rays $w_0^\pm$, $s^\pm(m_0)$, where $\angle_{o_0}(s^\pm(m_0), w_0^\pm) = 2m_0$, and $s^\pm(m_0) \cap F \neq \emptyset$.

Therefore, it follows from $\sum_{w \in A} \tilde{\psi}_w = \pi$ that $\sum_{w \in A_0} \tau_w \geq m_0$, which completes the proof. $\square$

3 Accumulating procedure

To describe the accumulating procedure we need some information about metric structure of the universal covering X of (M, g) , where g is a geometrization.

3.1 Metric structure of the universal covering

Recall (see, for example, [BS], [CK]) that X can be represented as the countable union $X = \cup_v X_v$ of blocks, where each X_v is a closed convex subset in X isometric to the metric product $F_v \times \mathbb{R}$, and F_v is the universal covering of a compact hyperbolic surface S_v with geodesic boundary. Every two blocks are either disjoint or intersect over a boundary component which is a 2-flat in X separating them and consequently no three blocks have a point in common. The 2-flats in X , which separate blocks, are called the *walls*. A wall w common for blocks $X_v, X_{v'}$ covers a 2-torus $e \subset M$, which separates (may be locally) the blocks $M_v = \pi(X_v)$, $M_{v'} = \pi(X_{v'})$ of M . The metric decompositions $X_v = F_v \times \mathbb{R}$, $X_{v'} = F_{v'} \times \mathbb{R}$ do not agree on w , and their $\mathbb{R}$ -factors induce two fibrations of w by parallel geodesics. We denote by α_w the angle between these fibrations, $0 < \alpha_w \leq \pi/2$. Since M is compact and the set E of separating tori in M is finite, we have $\alpha_0 := \inf_w \alpha_w > 0$, where the infimum is taken over all the walls in X .

3.2 Modified Poincaré series

We fix a wall $w^* \subset X$, take a block $X_{v^*} \subset X$, for which w^* is a boundary wall, and take a base point $x_0 \in w^*$. We denote by W_0 the set of the boundary walls of X_{v^*} different from w^* ; by W_n , $n \geq 1$ the set of walls in X at the combinatorial distance $n + 1$ from w^* , that is, $w \in W_n$ if and only if any geodesic segment in X between x_0 and w intersects n walls over its interior, including a wall from W_0 . Note that $W = \cup_{n \geq 0} W_n$ consists of all walls in X , which lie on one and the same side of w^* as X_{v^*} . Now, we define

a modified Poincaré series as follows

$$\mathcal{P}_W(t) = \sum_{w \in W} e^{-t \operatorname{dist}(x_0, w)}.$$

Comparing P_W with $\mathcal{P}(t) = \sum_{\gamma \in \Gamma} e^{-t|x_0 - \gamma x_0|}$, one easily obtains from triangle inequality that $\mathcal{P}(t) \geq e^{-D} \mathcal{P}_W(t)$, where $D > 0$ is the maximal diameter of the tori $e \in E$. Recall that the critical exponent of $\mathcal{P}$ is defined as the infimum of $t \in \mathbb{R}$ for which $\mathcal{P}(t) < \infty$. Therefore, the critical exponent $\bar{h}$ of $\mathcal{P}_W$ satisfies $\bar{h} \leq h(g)$, and to prove Theorem 1.1 it suffices to show that $\bar{h} > 1$.

3.3 Special broken-geodesic paths

Proposition 3.1. *For each $n \geq 0$ we have*

$$\mathcal{P}_n(1) := \sum_{w \in W_n} e^{-\operatorname{dist}(x_0, w)} \geq \frac{\pi}{4} \lambda_0^n,$$

where $\lambda_0 > 1$ is the constant from Lemma 2.1.

Proof. Induction over n . For each $n \geq 0$ and each wall $w \in W_n$ we produce a broken geodesic ξ_w in X between x_0 and w as follows. For $w \in W_0$ we put $\xi_w = x_0 x_w$, where $x_w \in w$ is the point closest to x_0 . Note that $x_0 x_w$ lies in a horizontal slice $F_{v^*} \times \{r_0\}$ of the block $X_{v^*} = F_{v^*} \times \mathbb{R}$.

Assume that for all k , $0 \leq k \leq n-1$ and all $w \in W_k$ the broken geodesic ξ_w is already defined, ξ_w is a juxtaposition $\eta_{w_0} \eta_{w_1} \dots \eta'_{w_k}$ of geodesic segments with the sequence $w_i \in W_i$ of walls leading to $w = w_k$. Moreover, we assume that each segment of ξ_w lies in a block and connects its different boundary components, the last segment η'_{w_k} lies in a horizontal slice of the block, which contains it, and η'_{w_k} is orthogonal to the wall w .

Take $w \in W_n$. Then there is a unique $\bar{w} \in W_{n-1}$ which precedes w . By the assumption, the last edge $\eta'_{\bar{w}} = s_{\bar{w}} x_{\bar{w}}$ of $\xi_{\bar{w}}$ lies in a horizontal slice $F_{v'} \times \{r_{v'}\}$ of its block $X_{v'} = F_{v'} \times \mathbb{R}$ and it is orthogonal to the wall $\bar{w}$ (at the end point $x_{\bar{w}}$). The important feature of our construction is that ξ_w contains all segments of $\xi_{\bar{w}}$ but the last one $\eta'_{\bar{w}}$.

Let $X_v \subset X$ be the other block adjacent to $\bar{w}$, in particular, the walls w , $\bar{w}$ are its boundary components. Recall that the metric splitting $X_v = F_v \times \mathbb{R}$ does not agree with that of $X_{v'}$ along of $\bar{w}$. Let $F_v \times \{r_v\}$ be the horizontal slice of X_v which contains $x_{\bar{w}}$ on the corresponding boundary component. The boundary lines of the slices $F_{v'} \times \{r_{v'}\}$, $F_v \times \{r_v\}$, which lie in $\bar{w}$, contain $x_{\bar{w}}$ and form an angle $\alpha_{\bar{w}} \in (\alpha_0, \pi/2]$.

To construct ξ_w we do the following. We take an isometric copy $F_v \subset \mathbb{H}^2 \times \{r_{v'}\}$ of $F_v \times \{r_v\}$ (by rotating the last by the angle $\alpha_{\bar{w}}$), where $F_{v'} \times \{r_{v'}\} \subset \mathbb{H}^2 \times \{r_{v'}\}$, so that $F_{v'} \times \{r_{v'}\}$ and F_v are sitting in the hyperbolic plane $\mathbb{H}^2 \times \{r_{v'}\}$ and they are adjacent along the common boundary

component, for which we use the same notation $\bar{w}$. Now, we connect the initial point $s_{\bar{w}}$ of the last edge $\eta'_{\bar{w}} \subset \xi_{\bar{w}}$ with the boundary component of F_v corresponding to w by the shortest geodesic segment $s_{\bar{w}}x'_w \subset H^2 \times \{r_{v'}\}$ and take $t_w = s_{\bar{w}}x'_w \cap \bar{w}$. The segment $t_w x'_w$ turned back to $F_v \times \{r_v\}$ gives the last segment $\eta'_w = s_w x_w$ of ξ_w . So $s_w \in \bar{w} \cap F_v \times \{r_v\}$, $|s_w - x_{\bar{w}}| = |t_w - x_{\bar{w}}|$ and $\eta'_w \subset F_v \times \{r_v\}$ is orthogonal to w at x_w . To complete the construction of ξ_w , we delete the last segment $\eta'_{\bar{w}} \subset \xi_{\bar{w}}$ replacing it by $\eta_{\bar{w}}\eta'_w$, where $\eta_{\bar{w}} = s_{\bar{w}}s_w \subset X_{v'}$. Clearly, so constructed ξ_w has all properties advertised above.

Let $l = |s_{\bar{w}} - x_{\bar{w}}| = L(\eta'_{\bar{w}})$ be the length of the last segment of $\xi_{\bar{w}}$, $l'_w = L(\eta_{\bar{w}})$, $l''_w = L(\eta'_w)$. Then we have

$$L(\xi_w) = L(\xi_{\bar{w}}) - l + l'_w + l''_w = L(\xi_{\bar{w}}) - l + \tilde{l}_w - \Delta_w,$$

where $\tilde{l}_w = L(s_{\bar{w}}x'_w)$, $\tilde{l}'_w = L(s_{\bar{w}}t_w)$ so that $\tilde{l}_w = \tilde{l}'_w + l''_w$, and $\Delta_w = \tilde{l}'_w - l'_w$. We obtained the same configuration which was studied in sect. 2, and we are going to apply Lemma 2.1 to estimate $\mathcal{P}_n(1)$ from below. Note that l being the length of a segment in X connecting different boundary components of a block is separated from 0 by some positive constant l_0 , which depends only on M , $l \geq l_0 > 0$.

For $n = 0$ we have

$$\mathcal{P}_0(1) = \sum_{w \in W_0} e^{-|x_0 - x_w|} = \sum_{w \in W_0} \tan \frac{\psi_w}{4} \geq \frac{\pi}{4},$$

where ψ_w is the angle under which the boundary component w of X_{v^*} is observed from x_0 (in the horizontal direction).

By inductive assumption we have

$$\mathcal{P}_{n-1}(1) \geq \sum_{w \in W_{n-1}} e^{-L(\xi_w)} \geq \frac{\pi}{4} \lambda_0^{n-1}.$$

Represent $W_n = \cup_{\bar{w}} W_{n, \bar{w}}$, where the union is taken over all $\bar{w} \in W_{n-1}$ and each $w \in W_{n, \bar{w}}$ follows $\bar{w}$. Then

$$\mathcal{P}_n(1) \geq \sum_{w \in W_n} e^{-L(\xi_w)} = \sum_{\bar{w} \in W_{n-1}} e^{-L(\xi_{\bar{w}})} e^l \sum_{w \in W_{n, \bar{w}}} e^{\Delta_w} e^{-\tilde{l}_w}.$$

Applying Lemma 2.1 with $A = W_{n, \bar{w}}$, we obtain

$$\mathcal{P}_n(1) \geq \lambda_0 \sum_{\bar{w} \in W_{n-1}} e^{-L(\xi_{\bar{w}})} \geq \frac{\pi}{4} \lambda_0^n,$$

which completes the proof. $\square$

4 Self-similarity argument

The constant $\lambda_0 > 1$ from Proposition 3.1 depends only on some metric data of M . Thus there is $n \in \mathbb{N}$, $n = n(M)$, so that $\frac{\pi}{4}\lambda_0^n > 1$. It follows from Proposition 3.1 that

$$\mathcal{P}_n(\bar{h}) = \sum_{w \in W_n} e^{-\bar{h} \text{dist}(x_0, w)} \geq 1$$

for some $\bar{h} > 1$. Furthermore, taking n sufficiently large, we can find $\bar{h} > 1$ with $\mathcal{P}_n(\bar{h}) \geq 1$ for any choice of the initial block X_{v^*} , its wall w^* and the base point $x_0 \in w^*$, since the set of choices up to isometries of X is compact.

We fix $n \in \mathbb{N}$ with this property and select a subset $W^* \subset W$, $W^* = \cup_{k \geq 1} W_k^*$, where $W_k^* = W_{kn}$. The set W_1^* serves as the generating set for W^* . Connecting x_0 with each wall $w \in W_1^*$ by the shortest geodesic segment $x_0 x_w$, we obtain new base points $x_w \in w$ (these x_w may be different from x_w constructed in the proof of Proposition 3.1). By induction, we find a base point $x_w \in w$ for each $w \in W_k^*$, $k \geq 1$ with the property $\text{dist}(x_{\bar{w}}, w) = |x_{\bar{w}} - x_w|$, where $\bar{w} \in W_{k-1}^*$ precedes w . Furthermore, by the choice of n and $\bar{h}$ we have

$$\sum_w e^{-\bar{h}|x_{\bar{w}} - x_w|} \geq 1,$$

for each $\bar{w} \in W_{k-1}^*$, where the summation is taken over all $w \in W_k^*$ which follow $\bar{w}$. Since $\text{dist}(x_0, w) \leq \text{dist}(x_0, x_w) \leq \text{dist}(x_0, x_{\bar{w}}) + |x_{\bar{w}} - x_w|$, we obtain

$$\begin{aligned} \mathcal{P}_{W_k^*}(\bar{h}) &= \sum_{w \in W_k^*} e^{-\bar{h} \text{dist}(x_0, w)} \\ &\geq \sum_{w \in W_{k-1}^*} e^{-\bar{h} \text{dist}(x_0, x_w)} \geq \sum_{w \in W_n} e^{-\bar{h} \text{dist}(x_0, x_w)} \geq 1 \end{aligned}$$

for each $k \geq 1$. Therefore, the modified Poincaré series

$$\mathcal{P}_W(\bar{h}) \geq \mathcal{P}_{W^*}(\bar{h}) = \sum_{k \geq 1} \mathcal{P}_{W_k^*}(\bar{h})$$

diverges at $\bar{h}$ and hence $h(g) \geq \bar{h} > 1$. This completes the proof of Theorem 1.1.

References

- [BW] W. Ballmann and M. Wojtkowski, An estimate for the measure theoretic entropy of geodesic flow, *Ergod. Th. & Dynam. Sys.* 9 (1989), 271–279.

- [BK] S. Buyalo and V. Kobel'skii, Geometrization of graph-manifolds. II. Isometric geometrization, *St. Petersburg Math. J.* 7 (1996), 387–404.
- [BS] S. Buyalo and V. Schroeder, On the asymptotic geometry of nonpositively curved graph manifolds, *Trans. Amer. Math. Soc.* 353 (2001), 853–875.
- [CK] C. Croke and B. Kleiner, The geodesic flow and a nonpositively curved graph manifold, *arXiv:math.DG/9911170*, 1999.
- [HS] C. Hummel and V. Schroeder, Tits geometry of cocompact real-analytic Hadamard manifolds of dimension 4. *Differential Geom. Appl.* 11 (1999), 129–143.
- [L] B. Leeb, 3-manifolds with(out) metrics of nonpositive curvature. *Invent. Math.* 122 (1995), 277–289.
- [M] A. Manning, Topological entropy for geodesic flows. *Ann. of Math.* (2) 110 (1979), 567–573.